



HIV Counseling and Testing Uptake, Knowledge, Attitude and Influencing Factors among Student Nurses and Midwives in the Gambia: An Institutional -Based Cross-Sectional Study

Sang Mendy^{1,5}; Ousman Bajinka^{2,3,5}; Amadou Barrow^{4,5}; Sun Mei¹

¹Xiangya School of Nursing, Central South University (CSU), Changsha, Hunan Province, China.

²Department of Microbiology, Central South University Changsha, Hunan Provinces, China.

³China-Africa Research Centre of Infectious Diseases, School of Basic Medical Sciences, Central South University, Changsha 410078, Hunan, China.

⁴Heidelberg Institute of Global Health, University Hospital and Medical Faculty, Heidelberg University, Heidelberg, Germany.

⁵School of Medicine and Allied Health Sciences, University of the Gambia, Banjul, the Gambia .

*Corresponding Author(s): Sun Mei

Xiangya School of Nursing, Central South University,
Changsha 410078, Hunan, China.

Fax: 86-13787793984; Email: sunmei@csu.edu.cn

Received: Dec 08, 2022

Accepted: Dec 27, 2022

Published Online: Dec 29, 2022

Journal: Annals of Epidemiology and Public Health

Publisher: MedDocs Publishers LLC

Online edition: <http://meddocsonline.org/>

Copyright: © Mei S (2022). *This Article is distributed under the terms of Creative*

Commons Attribution 4.0 International License

Keywords: Barriers; Facilitators; HIV testing and counseling; Student nurses; Uptake & attitudes.

Abstract

Introduction: The key intervention in prevention and controlling the spread of HIV, counseling and testing (HCT) is pertinent across the globe. Despite this pandemic, However, HCT intervention is still undertaken and more especially, in sub-Saharan Africa. Student nurses formed the largest group undergoing health care training in the country compared to doctors and other health cadres. Also, student nurses are usually in their youthful age and according to WHO, they are part of the most vulnerable group to HIV infection. They are also expected to provide HIV services upon graduation, judging by the fact that they interact more with patients/clients than other healthcare professionals. In response to this, we designed this study to determine the prevalence of HCT uptake, knowledge, and attitude and further evaluated the factors that influence the student nurses and midwives in public nursing schools.

Methods: Our design involved an institutional-based cross-sectional study to collect data from 305 nursing students and midwives selected randomly with the use of validated and reliable self-administered questionnaire. SPSS version 25.0 was used for descriptive statistics (percentages, mean and standard deviation) and inferential statistics (chi square, logistics regression, one way ANOVA) and independent samples t-test. The statistical significance was considered at p-value <0.05.



Cite this article: Mendy S, Bajinka O, Barrow A, Mei S. HIV Counseling and Testing Uptake, Knowledge, Attitude and Influencing Factors among Student Nurses and Midwives in the Gambia: An Institutional-Based Cross-Sectional Study. *A Epidemiol Public Health*. 2022; 5(2): 1100.

Results: Out of the 305 students recruited for the study, 60.98% were females, with a mean age of 25.5 years old. Our study found that 58.4% of the participants had tested for HIV in the past. When compared to the younger age between 19 and 24, the prevalence of HCT utilization or uptake was higher among the age 31 and above. At least 67% of the participants with positive attitude towards HTC are more likely to utilize HCT against those with negative attitudes [OR= 1.67, 95% CI: (1.009–2.76), $p= 0.046$]. With regards to those who will undergo HIV testing, we found that age group 25 to 30 years are less likely when compared with those within the age range of 19–24 years [OR =0.282, 95% CI: (0.14-0.567), $p<0.001$]. Department category to utilize HCT, we found that 2.7 times more likely is from midwifery students when compared with third year nursing students [OR= 2.718, 95% CI: (1.124-6.575), $p=0.027$]. Finally, a remarkable number (95.7%) acknowledged the benefits attached to HCT in preventing and controlling the spread of HIV/AIDS pandemic.

Conclusions: HCT uptake among student nurses and midwives was influenced by factors ranging from individual to interpersonal challenges (such as concerns of friends), perceived susceptibility to the disease, and lack of confidentiality, stigma and discrimination from health service providers. Therefore, these barriers can be addressed through an organized targeted health education intervention and advocacy programs across health training institutions in the Gambia and beyond.

Introduction

Human immunodeficiency virus (HIV) infection is now predominantly seen as a controllable condition due to treatment advances and care. However, without appropriate strategy or intervention, HIV can lead to Acquired Immune Deficiency Syndrome (AIDS), which may bring about morbidity, health burden and mortality [1]. Mitigating the rapid transmission of HIV requires concerted efforts and commitment. This involves widespread screening and testing to isolate persons who may miss opportunities to seek treatment and thereby unknowingly transmitting the virus to others. It is apparent that HIV counseling and testing (HCT) is a key strategy in curbing the spread of HIV [1]. However, this is only achievable with attitudinal change, and a low frequency in the uptake of HIV counseling and testing is a challenging issue among healthcare provider [2]. Great awareness and optimistic perception of HCT health professionals are essential prerequisites [3]. From the literature, HIV/AIDS and attitude towards HCT are effective among health workers including pre-service nurses [4,5]. Thus, student nurses and midwives with good attitude towards HCT will educate clients [6].

Globally, about half HIV/AIDS patients become infected before age 25 years, making HIV as the second most common cause of death between the age group 20 and 24 [7]. Student nurses fit well in this population cohort as they are usually fresh secondary school leavers with little guidance from their parents during the period of study. Student nurses and midwives are part of this vulnerable group that often contact their peers. Also, student nurses formed the largest group undergoing health care training in the country compared to doctors and other health cadres. Also, student nurses are usually in their

youthful age and according to WHO, they are part of the most vulnerable group to HIV infection. Student nurses are also expected to provide HIV services upon graduation, judging by the fact that they interact more with patients/clients than other healthcare professionals. Furthermore, the 2013 National Demographic Health Survey (DHS) found a significantly high HIV prevalence among youth aged 15–24 at 0.3% [8]. This figure includes student nurses, hence, given this tendency, preventive interventions that can limit the epidemic among the cream of our society and the larger population are crucial.

Despite the acceptance of the HCT model by African governments, the level of acceptance and utilization of this model by citizens has been met with mixed attitude [9]. Until today, the findings from studies conducted in Africa, regarding HCT uptake come with high efficacy acceptance [10]. Cost-effective treatments offer chances to raise awareness of HIV and preventative practices and these are key components of HIV prevention programs [11]. Knowing one's HIV status is crucial for prevention and counseling can be an important prevention and care strategy [12]. Community awareness was found to be associated with significant uptake of HCT in the women population who received pre-test counseling [13]. However, little is known about the level of knowledge on HCT among student nurses and midwives in The Gambia. Notwithstanding, the 2013 national demographical health survey revealed that knowledge of places for HIV testing is higher among urban women and men than in rural areas. Moreover, despite high knowledge of the sources of HIV testing, only 39% and 19% of women men respectively had been tested at least once [14]. Also, Multiple Indicator Cluster Survey (MICS-6) report, 2018 indicated that 62.5% of men who knew a place to get tested, only 23.3% were tested and 22.4% of men had been tested and knew the most recent test result, and in the past 12 months, only 8.0% had been tested and are conscious of their status [7]. Apparently, a study found that those with more negative attitude towards HIV are did not undergo HIV test [15].

Globally, numerous studies have outlined many barriers and facilitators that can influence HIV counselling and testing in different population. Barriers and facilitators may also differ across countries and group of people with different characteristics. Findings from many studies also grouped barriers related to HCT into five main domains based on the socio-ecological model (interpersonal, intrapersonal, community, institutional and policy levels). In other studies, the barriers are phrased differently but meaning the same. For example, the intent to seek for HIV counselling and testing was attributed or related to five attitude subscales namely: people's concerns, individual concerns, friends' concerns, value attached to testing, and confidentiality, perceived susceptibility [18]. Therefore, the magnitude of each of these independent domain could be unevenly distributed at national level and globally. However, the factors that influence HCT uptake among student nurses and midwives in the Gambia are not known and it is obvious that this study hope to determine these factors. The study also examined utilization rate or prevalence, knowledge and attitude towards HCT uptake. Findings expected to be identified from this will enable policy makers to redesign some of the policies and regulations on HIV/AIDS and HCT services at national level. Hence, these findings may trigger a need to overhaul some policies or regulations that impedes HCT uptake in the country.

Methods

Design and setting

In our study, we conducted a cross-sectional study for the three public nursing schools in different regions. These are: Registered School of Nursing and Midwifery at Banjul (capital city); School of Enrolled Nursing and Midwifery (Bansang town) and and the Community health and midwifery (Mansakonko town). The School of Enrolled Nursing and Midwifery is located in Central River Region in the rural part of the country. It is about 300km from the capital Banjul and it has a two years nursing program and one year midwifery program with residence or accommodation available for students.

Similarly, the School of Community Health and Midwifery is located in Lower River Region about 200km from the capital. It also has a two year program and a one year midwifery program. Unlike the other two schools, graduates are usually posted to the community and less often to minor health centres where they engage in community. Students in this school are not provided with residence or accommodation, hence are forced to rent within the vicinity of the school and sometimes very far away from the school where lecture are conducted.

The School of State Registered Nursing and Midwifery is located in Banjul under the Gambia College is a three year registered nursing program and 18 months diploma midwifery program. It has a more comprehensive nursing curriculum compared to the other two schools. It also accommodate some of its students within the school campus while majority stay outside.

Sample and Eligibility Criteria

Our study population included student nurses and midwives who are currently studying in the three public nursing schools in The Gambia. They are usually in their youthful age and according to WHO, they are part of the most vulnerable group to HIV infection. They are also expected to provide HIV services upon graduation judging by the fact that they interact more with patients/clients compared to other health care professionals.

Student nurses and midwives presently studying in public nursing schools and willing to participate were recruited. However, the study excluded support staff, lecturers and practicing nurses.

Sample size determination

In other to determine the sample size for the study, we used Yamane (1967) formula to calculate the number of student nurses and midwives to participate in the study. The formula, assuming a confidence interval of 95% and error of 5%, with a population of 651 students from all three schools, yielded a desired sample size of 248 participants. However, 10% was added to cover for the non-response rate and erroneous questionnaires after completion; hence, the final desired sample size was 273. However, a total of 305 students volunteered to participate surpassing he desired calculated sample size for the study. Below is a breakdown of how the researcher arrive to the sample size for the study.

$$n = \frac{N}{1+N(e)^2}$$


Where n= the sample size N = the student population e= the error of 5% points (0.05)

We added 10% to cover for non-response rate and erroneous questionnaires after completion thus, final desired sample size was 273.

Sampling technique

Participations were selected using stratified random sampling technique from the three schools. An enumerated list of all the regular students enable the study have a clear idea as to the number of active students for each of the academic years and used as sampling frame. Proportional allocation was done for each of the academic levels i.e. year 1, 2, 3 (RN school only) and midwifery class. Stratified random sampling using lottery method was used to draw samples from each of the respective academic levels or classes. A total of 305 students volunteered to participate in the study.

Schematic presentation of the sampling method



Measurement

A self-administered questionnaire segmented into three parts was used to collect data from participants. The segments comprised of the following parts: The socio-demographic information which includes, sex, age, marital status, level of education of parents, religious affiliation, place of residence; ethnicity, school, academic level, marital status, educational level of parents, wealth quintile, and Local Government Area (LGA) of the participants. The variables used herein are from previous published studies and also demographic health survey of the Gambia, 2018.

The items on knowledge are subset of items provided by the World Health Organization as part of its research guidelines for studies related to HIV counseling and testing [19]. This section contained questions about knowledge on HCT such as: source of information about HCT, perception on the benefits of HCT, practice of HCT, satisfaction with HCT services, reason for not taking HCT, willingness to HCT and preference of HCT method. The section contained 24 items with responses of yes/no and also multiple choices. Analysis was done based on the individual items mentioned above using descriptive analysis in the form of frequency, percentages and means.

For attitude, 32-item HIV-Antibody Testing Scale was used (Boshamer CB, Bruce KE)[20]. This scale was validated and reliable as it was used in different previous studies in Africa and beyond with a Cronbach's alpha of 0.88. Responses were scored on the basis of strong agreement (given 5) and strong disagreement (given 1). An attitude score of more than or equals to the

total mean score was regarded as positive attitude and less than the total mean score was considered as negative attitude [21]. The test-retest reliability result for the HIV-Antibody Testing Scale showed a good internal consistency with a Cronbach's alpha score of 0.813 for the present study. Equally, the knowledge questionnaire part had a good internal consistency with a Cronbach's alpha of 0.783.

Data collection

We used self-administered questionnaire to collect data and the assistance of data collectors in each school upon receiving permission from the heads of schools. All the data collectors were trained prior to data collection. Data collectors in each school informed all students about the research prior to the data collection process. Student representatives were also used to inform their colleagues on the importance and benefits of the research. The time for data collection was communicated to all the lecturers and students in the three nursing institutions.

Statistical Analysis

We used Statistical Package of Social Sciences (SPSS) version 25 software for data entry and statistical analysis. Descriptive and frequency analyses were used to analyze socio-demographic characteristics of the participants, knowledge items and independent variables respectively. Categorical variables were presented as frequencies and percentages and chi square test was used to test for association. If the number of participants in one or more categories was less than five, Fisher's exact test was applied. Results were considered to be significant with a p -value < 0.05 . Multivariate logistic regression was used to investigate the predictors or demographic variables association with HCT uptake and conditional method calculating odds ratio and 95% confidence interval was applied. One-way ANOVA and independent samples t-test were used to determine the mean difference between categories of the demographic variables and dependent outcomes for the attitude scale.

Factor analysis was also conducted to identify barriers and facilitators associated with HCT among student nurses and midwives using the 32 items on the HIV Antibody Testing Scale. A principal component analysis was used for facilitators and barriers using varimax rotation together with scree plot for the extraction of factors. We applied a cut-off point of 0.40 for item inclusion in the interpretable factors and also the items with two or more loadings >0.30 were considered cross loading items. We assigned to a single factor with the highest loading [22]. Furthermore, eigenvalue of more than 1 points was taken into consideration in identifying factors using the scree plot. All the items loaded above 0.40 cut-off.

Ethical considerations

Prior to the study, the study was granted ethical approval from the Institute Review Board at Central South University, Xiangya School of Nursing, China (IRB Number: E202044). Similarly, approval to conduct the research in the nursing schools was sought and approved by the Gambia Scientific Coordination Committee (GSCC) of the University of the Gambia. Confidentiality was maintained and the participation to this study was strictly on voluntary. Furthermore, no incentives were provided for participation and the participants were made to understand that they could withdraw from the study at their will at any time or stage of the study. Also, respondents were issued with

a detailed information sheet about the study and consent form was signed by participants willing to participate. Confidentiality was maintained and participants were not required to write any information that may reveal their identity. Participation was voluntary, and no incentives were provided for participation. Apart from time, there was no other potential risks related to the study participants or the information they provided. Furthermore, this study was not invasive therefore, no harm was anticipated on the participants. There was no repercussion or punishment to be meted on the students for declining to participate in the study.

Results

Socio-demographic characteristics of the participants

In this study, the response rate was 91% (305 submitted questionnaires out of 335 were accepted for analysis) and out of the 305 participants, more than half (60.9%) were females. For the mean age and the standard deviation, nursing and midwife students was 25.5 years \pm 5.4. However, by age range, more than half of participants were within age group of 19-24, at 58.3%. HCT use was more widespread within the age range of 31 & above years at 80.0% compared to 19-24 years at 43.5%. The HIV counseling and testing uptake among midwifery students was also higher by 81.5% compared to the 58.8% second-year students. Students from Janjanbureh as well as Mansakonko LGA chose HIV counseling more than other LGAs in the study. These LGAs are located in rural areas of the country, which are frequently plagued by a lack of resources. The Mandingo tribe was the most common ethnic affiliation of the participants at 40.9%, while about 92.1% were Muslims. In terms of distribution of participants by school, about 43.6% were from Registered Nursing School of Gambia College, 31.8% from Community Health Nursing School, Mansakonko and 24.6% from State Enrolled Nursing, Bansang. The majority of participants were single at 65.3% while almost one in every four students were married at the time of the study. About 36.1% were in their second academic year as shown in **Table 1**. Regarding the educational status of participant parents, almost half of participants' father and two-third of their mothers had no education. Our participants' wealth quintile was largely in the middle index class at 48.5%, followed by the second wealth index at 25.3%. Approximately about half of the participants report to be living with both parents while about one-third report to have only one parent alive and one in five reported to have lost all their parents. According to participants' LGA local of origin, Brikama has the most residents while Kuntaur had the lowest participants at 1.7%. In overall, about 57.1% were from urban areas of the country.

HIV counselling and testing uptake

HIV counseling and testing uptake among nursing students and midwives was 58.4% as shown in **Figure 1**. HCT utilization was more prevalent within the age range of 31 & above years, 32 (80%) compared to the 19-24 years of age 77 (43.5%). Also, HIV counseling and testing utilization was higher among the midwifery students, 53(81.54) compared to second year students, 87(58.78). Students who are from Local Government Area (LGA) of Janjanbureh opted for HIV counseling more than other LGAs, 18(81.82%) followed by Mansakonko LGA, 19(79.17). These LGAs are situated in the rural part of the country which are often plagued by limited resources (**Table 1**).




Figure 1: HIV counselling and testing uptake.

Knowledge on HCT

Regarding source of information about HCT, majority of the respondents heard of HCT before 290(95.08%). The most common source of information was at school by 53.1% and followed by health facility at 34.4% as shown in Figure 2. Of the 198 participants who had the HIV test, the very common reason for having done the test was on the interest of aware of self-status scored 65.3% while for blood donation was at 23.2%. Those of satisfied with HCT gave a score of 95.5%. Almost all the participants (99.7%) agreed to the fact that HIV counseling and testing is important for the control of HIV/AIDS whilst 87.2% believed HCT is beneficial to both positive and negative persons. About 75.4% of the participant indicated that they will prefer confidential-linked testing method compared to anonymous /self-testing. Participants were also asked as to who should go for HIV testing, 67.5% mentioned anyone at risk and 23.0% mentioned only for those who are sexual active with multiple partners and about two-third preferred to receive counseling from a trained counselor compared to a nurse or doctor.

Despite many participants not having a HIV test in the past, 94.8% reported their desire to go for a test in the future. In terms of participants' health facility of preference to have a HIV test, 67.8 % mentioned government health institutions having HCT services followed by preference for private health facilities/clinics at 24.6%.




Figure 2: Source of information about HIV counseling and testing among respondents.

Attitude towards HCT

The mean attitude score was 113.94±12.23. A total of 149(48.9%) of the students showed negative attitude towards HCT based on the overall mean attitude score. The results also showed that out of the students found in State Registered Nursing School, at least half (52.6%) of them showed a negative attitude towards HCT when compared to with others; community health nursing school (46.6%) and state enrolled nursing school (43.3%).

Majority of the students agreed (94.1%) that it is very im-

portant to seek for HIV and 86.9% rejected the statement: "HIV testing is unnecessary for me, as I feel healthy". However, about one-third of the participants assumed that everyone who is tested is infected with HIV, while 19.7% stayed neutral about this assumption (Table 2). Further analysis revealed the same trends in participants' opinions on HIV test information being kept very confidential by the medical staff who do the testing, the tendency to be judged or ignored by friends who knew they had done the test, and a very low proportion of neutral stands on this item. Additionally, more than half of the students disagree with the possibility of being positive for HIV/AIDS unknown to them, as well as having intercourse with a person who was at risk for HIV/AIDS.

Socio-demographic variables associated with HCT uptake

To explore the association between demographic variables and HCT uptake, chi square analysis was conducted. The analysis revealed statistically significant association between six independent variables (age, religion, marital status, academic year of study, academic class by school and LGA) and HIV uptake as shown in Table 1. Age was statistically significant ($\chi^2=38.342, p<0.001$), religion ($\chi^2=9.136, p=0.003$), academic class of study participants ($\chi^2=21.628, p<0.001$), marital status ($\chi^2=42.110, p<0.001$) and year of study by school ($\chi^2=27.887, p=0.001$) and LGA ($\chi^2=19.340, p=0.003$) (Table 1). Other influencing factors associated with HIV counselling and testing were attitude ($\chi^2=7.720, p=0.005$), heard of HCT before ($\chi^2=6.521, p=0.011$), HCT availability ($\chi^2=8.336, p=0.004$) and knowing where to get HCT services ($\chi^2=12.283, p<0.001$).

Four variables showed statistically significant relationship with HCT uptake. The model was statistically significant using forced entry method ($\chi^2=71.219, p<0.001$) which clearly showed its capability of detecting between participants who utilized HCT and those who did not. The overall explanation of the model was 28% (Cox & Snell R²) and 20.8% (Nagel Kerke R²) of the variance in the utilization of HCT.

In comparing the determinants of HCT uptake among the participants, the analysis showed Christians were five times more likely not to utilize HCT services compared to Muslim participants [OR = 5.272, 95% CI (1.839-15.112), $p=0.002$]. Similarly, participants who are single are four times more likely not to undergo HIV counselling compared to married participants [OR =4.265, 95% CI (1.866-9.746), $p=0.001$]. Also, participants who are in a relationship where seven times more likely not to take a HIV test compared to married students [OR=7.361, 95% CI (1.768-30.656), $P=0.006$]. Furthermore, participants within the age range of 25-30 years are more likely to undergo HIV testing than those within the age range of 19-24 years [OR =0.282, 95% CI: (0.14-0.567), $p<0.001$]. In academic class comparison, mid-wifery students were 2 times more likely to utilize HCT [OR=2.718, 95% CI (1.124-6.575), $p=0.027$] compared to third year students as shown in Table 3.

Influencing factors associated with HCT uptake

We used the HIV Testing Antibody Scale (HTAS) with 32 items to identify factors influencing HIV counselling and testing uptake. For the overall scale, the data was adequate for factor analysis with an overall Kaiser-Meyer-Olkin Measure of Sampling Adequacy (KMO) of 0.759. The instrument was further sectioned into two subscales of facilitators and barriers associated with HIV counselling testing. The facilitators were 17 items and barriers are 15 items in total. To identify facilitators

and barriers associated with HIV counseling and testing among nursing students, factor analysis we used principal component and Varimax rotation for the different subscales.

Facilitators to HIV counseling and testing

The items recommended as facilitators were optimal for factor analysis (KMO = 0.685). A total of five factors with an Eigen value more than 1 were identified using the Scree test. These factors gave a total of 54.7% of the variance in scores. These identified factors and their loadings are tabulated in Table 4.

Factor one (Eigen value = 3.09) gave 18.2% variance in the responses, which is largely about the support from friends or partners in relation HCT uptake. The factor comprised of items about the concerns of friends' feeling/reactions or support towards HIV counseling and testing. Items such as: "my friends would not treat me any differently if I were tested for HIV" and "my friends would not look down on me if I were tested for HIV" loaded highly in this factor (Table 5). A total of five items aligned under this factor accruing the majority of the items under this subscale. Support and positive attitude of friends is an important facilitator associated with HIV counselling and testing among nursing students. Therefore, the attitude of peers, partners or family members can immensely facilitate the uptake of HCT among nursing students.

The second factor (eigenvalue = 2.02) accounted for 11.86% of the variance and is associated with the intrapersonal level factors under the socio-ecological model. It is mainly concerned with the individual's perceived susceptibility to HIV/AIDS infection and the need for HCT. Self-evaluation is a crucial ingredient in promoting the utilization of HCT supported by the cross-cutting factors like knowledge, motivation and ability to act or take a decision. The following items loaded highly; 'There is a possibility that I have HIV and AIDS', 'I may have had sex with someone who was at risk for HIV/AIDS' and 'I am at risk for HIV [Table 4].

The third factor (eigenvalue = 1.69) with an overall variance of 9.96% is related to personal concerns about HCT as a whole. The value of HCT, attitude of staff and family members are key to facilitating the uptake of HCT. The utilization of HCT will be enhanced if these sub-factors are address at our health facilities where services are rendered. Items that loaded high included; 'HIV tests give accurate results', 'I would be comfortable talking to an HIV counselor about personal behaviors that place me at risk for HIV infection', 'It is extremely useful to test for HIV' and 'I could easily discuss HIV-antibody testing with my family'. The clearly demonstrated that if a good reception is accorded to clients at the health institution they will be more than willing to discuss further about HCT with family members thereby increasing the uptake as shown in **Table 4**.

Factor four (eigenvalue = 1.85) accounted for 7.56% variance in the responses and mainly comprised of items concerned with confidentiality and privacy. The items that loaded more with this factor are: 'I trust the HIV test counselors and nurses to keep my information confidential'. 'HIV test information is kept very confidential by the medical staff who do the testing'. 'Testing and counselling is a pleasant experience'. Clients would be motivated to utilize HCT if their privacy and confidentiality is assured by health staff and family members (**Table 4**).

The fifth factor (eigenvalue = 1.21) gave 7.13% of the variance in the response and is mainly concerned about the apprehension/fear and stigma associated with a HCT and a possible positive result. The items that loaded strongly with the factor are: 'I consider going for HIV counselling and testing extremely frightening' and 'I would like to be alone when doing the test' (**Table 5**). Demystifying the fear factor could be a great motivator for the uptake of HCT among nursing students and the general population at large.

Barriers to HIV counseling and testing

Similarly, the barrier items in the HIV testing Antibody scale were also applicable for factor analysis (KMO = 0.810) and five factors with an Eigen value more than 1 were identified using the Scree test. The factors accounted for a total of 59.21% of the variance in scores. The identified factors and their loadings are tabulated in **Table 5**.

The first barrier factor associated with HCT utilization had an Eigen value of 3.96 and gave 26.43% of the variance in the participants' responses. A total of five items aligned with this factor. The items contained in this factor are mostly about personal concerns and about peers/friends, partners' reaction to HIV counseling and testing. Five items loaded with this factor and item: 'I would not get tested for HIV because I would be asked information that was too personal' scored highly followed by the item "I do not have time to get an HIV test". Lack of support or negative stereotype from peers or friends can adversely deter individuals from HCT services as shown in **Table 5**. The second barrier factor with an eigenvalue of 1.58 and total response variance of 10.53% and a total of four items loaded strongly with this factor. The factor is mainly concerned with the value or perception attached to HCT by clients. Items such as: 'Anyone who is tested for HIV is dirty' and "admitting that you should be tested for HIV means that you have engaged in immoral behavior" scored highest followed by the item; 'HIV testing is unnecessary for me, as I feel healthy' as indicated in **Table 5**.

The third barrier factor (eigenvalue 1.28) had an overall response variance of 8.51% in which three items loaded strongly with the factor. The three items that aligned with this factor are mostly concerned about concerns of people with regards to HCT. Items such as 'people assume that everyone who is tested for HIV is infected with HIV', 'people would assume I have HIV if I decided to get tested' and 'I am afraid that if I were tested for HIV, my name would go into public record'. The fourth barrier factor (eigenvalue =1.06) had an overall response variance of 7.06% in which only two items loaded strongly. The two items that aligned with this factor are concerned about "*privacy & stigma associated with HCT and attitude of staff*". The two items are: "I would not want anyone to know if I got an HIV test" and "It would be embarrassing to get tested for HIV".

The fifth barrier factor (eigenvalue = 1.002) had an overall response variance of 6.68% and only one item loaded strongly with the factor which is *concerned about confidentiality (HIV-antibody testing is not really confidential)*.

Table 1: Socio-demographic characteristics and utilization of HCT (n=305).

Variable	Frequency n (%)	Ever had HIV test		Chi-square test	p-value
		Yes n (%)	No n (%)		
Age				38.342	0.001*
19-24 years	177 (58.03)	77 (43.50)	100 (56.50)		
25-30 years	88 (28.85)	69 (78.41)	19 (21.59)		
30 & above	40 (13.11)	32 (80.00)	8 (20.00)		
Gender				0.136	0.712
Male	119 (39.02)	71 (59.66)	48 (40.34)		
Female	186 (60.98)	107 (57.53)	79 (42.47)		
Religion				9.136	0.003*
Islam	281 (92.13)	171 (60.85)	110 (39.15)		
Christianity	24 (7.87)	7 (29.17)	17 (70.83)		
Ethnicity				13.361†	0.090
Mandinka	125 (40.98)	74 (59.20)	51 (40.80)		
Fula	68 (22.30)	40 (58.82)	28 (41.18)		
Wollof	30 (9.84)	19 (63.33)	11 (36.67)		
Jola	36 (11.80)	23 (63.89)	13 (36.11)		
Manjago	14 (4.59)	4 (28.57)	10 (71.43)		
Serere	13 (4.26)	8 (61.54)	5 (38.46)		
Aku	7 (2.30)	5 (71.43)	2 (28.57)		
Foreign	5 (1.64)	0 (0.00)	5 (100.00)		
Marital status				42.110	0.001*
Married	88 (28.85)	75 (85.23)	13 (14.77)		
Single	199 (65.25)	94 (47.24)	105 (52.76)		
Divorced	5 (1.64)	4 (80.00)	1 (20.00)		
Relationship	13 (4.26)	5 (38.46)	8 (61.54)		
Academic class				21.628	0.001*
First year	88 (28.85)	40 (45.45)	48 (54.55)		
Second year	110 (36.07)	59 (53.64)	51 (46.36)		
Third year	42 (13.77)	26 (61.90)	16 (38.10)		
Midwifery	65 (21.31)	53 (81.54)	12 (18.46)		
Year Study				27.887	0.001*
3rd Year RN	42 (13.8)	16 (38.1)	26 (61.9)		
2nd Year RN	32 (10.5)	17 (53.1)	15 (46.9)		
1st Year RN	33 (10.8)	22 (66.7)	11 (33.3)		
2nd Year CHN	44 (14.4)	19 (43.2)	25 (56.8)		
Midwifery SEN	14 (4.6)	1 (7.1)	13 (92.9)		
1st Year SEN	26 (8.5)	14 (53.8)	12 (46.2)		
2nd Year SEN	35 (11.5)	15 (42.9)	20 (57.1)		
Midwifery RN	26 (8.5)	4 (15.4)	22 (84.6)		
1st Year CHN	29 (9.5)	12 (41.4)	17 (58.6)		
Midwifery CHN	24 (7.9)	7 (29.2)	17 (70.8)		
Residence				0.693	0.405
Urban	174 (57.05)	80 (45.98)	51 (29.31)		
Rural	131 (42.95)	98 (74.81)	76 (58.02)		
Schools				0.731	0.694
SEN School	75 (24.59)	45 (60.00)	30 (40.00)		
CHN School	97 (31.80)	59 (60.82)	38 (39.18)		
RN School	133 (43.61)	74 (55.64)	59 (44.36)		
Parents alive				3.162	0.367
Yes	154 (50.49)	94 (61.04)	60 (38.96)		
Divorced	23 (7.54)	10 (43.48)	13 (56.52)		
One of them alive	108 (35.41)	64 (59.26)	44 (40.74)		
Both of them not alive	20 (6.56)	10 (50.00)	10 (50.00)		
Family type				0.143	0.705

Nuclear	157 (51.48)	90 (57.32)	67 (42.68)		
Extended	148 (48.52)	88 (59.46)	60 (40.54)		
Household wealth index				2.741†	0.611
Lowest	67 (21.97)	34 (50.75)	33 (49.25)		
Second	77 (25.25)	49 (63.64)	28 (36.36)		
Middle	148 (48.52)	87 (58.78)	61 (41.22)		
Fourth	7 (2.30)	4 (57.14)	3 (42.86)		
Highest	6 (1.97)	4 (66.67)	2 (33.33)		
LGA of origin				19.340†	0.006*
Kanifing	91 (29.84)	51 (56.04)	40 (43.96)		
Banjul	18 (5.90)	7 (38.89)	11 (61.11)		
Brikama	107 (35.08)	59 (55.14)	48 (44.86)		
Kuntaur	6 (1.97)	6 (100.00)	0 (0.00)		
Janjanbureh	22 (7.21)	18 (81.82)	4 (18.18)		
Mansankonko	24 (7.87)	19 (79.17)	5 (20.83)		
Kerewan	15 (4.92)	6 (40.00)	9 (60.00)		
Basse	22 (7.21)	12 (54.55)	10 (45.45)		

*Significant at p<0.05, †Fisher’s Exact test used

Table 2: Frequency analysis of items on attitude among nursing students (n=305).

Items	Strongly Agree/ Agree n(%)	Neutral n(%)	Strongly disagree/ Disagree n(%)
HIV-antibody testing is not really confidential	67 (22.0)	54 (17.7)	184 (60.3)
I would not want anyone to know if I got an HIV test.	148 (48.5)	60 (19.7)	97 (31.8)
HIV testing is unnecessary for me, as I feel healthy	21 (6.9)	19 (6.2)	265 (86.9)
I consider going for HIV counselling and testing extremely humiliating	39 (12.8)	28 (9.2)	238 (78.0)
People assume that everyone who is tested for HIV is infected with HIV.	133 (43.6)	27 (8.9)	145 (47.5)
Admitting that you should be tested for HIV means that you have engaged in immoral behavior	27 (8.9)	17 (5.6)	261 (85.6)
I am afraid that if I were tested for HIV, my name would go into public record	67 (22.0)	31 (10.2)	207 (67.9)
Anyone who is tested for HIV is dirty.	8 (2.6)	6 (2.0)	291 (95.4)
It would be embarrassing to get tested for HIV	25 (8.2)	20 (6.6)	260 (85.2)
People would assume I have HIV if I decided to get tested	106 (34.8)	60 (19.7)	139 (45.6)
I am afraid someone would find out I was tested for HIV	85 (27.9)	53 (17.4)	167 (54.8)
I would be embarrassed if my friends found out that I had decided to have HIV test.	61 (20.0)	45 (14.8)	199 (65.2)
I would not get tested for HIV because I would be asked information that was too personal	41 (13.4)	35 (11.5)	229 (75.1)
I do not have time to get an HIV test	24 (7.9)	19 (6.2)	262 (85.9)
My friends would treat me badly if I were tested for HIV	59 (19.3)	69 (22.6)	177 (58.0)
HIV test information is kept very confidential by the medical staff who do the testing.	49 (16.1)	40 (13.1)	216 (70.8)
My friends would not look down on me if I were tested for HIV	135 (44.3)	54 (17.7)	116 (38.0)
My friends would support my decision to get an HIV test	37 (12.1)	69 (22.6)	199 (65.2)
HIV tests give accurate results	28 (9.2)	51 (16.7)	226 (74.1)
It is extremely useful to test for HIV	15 (4.9)	3 (1.0)	287 (94.1)
I would be comfortable talking to an HIV counselor about personal behaviors that place me at risk for HIV infection	31 (10.2)	18 (5.9)	256 (83.9)
My friends would look down on me if I were tested for HIV	113 (37.0)	69 (22.6)	123 (40.3)
My friends would not treat me any differently if I were tested for HIV.	121 (39.7)	80 (26.2)	104 (34.1)
I would like to be alone when doing the test	17 (5.6)	12 (3.9)	276 (90.5)
I trust the HIV test counselors and 1s to keep my information confidential.	27 (8.9)	39 (12.8)	239 (78.4)
It would not bother me if someone I know sees me going to get an HIV test	107 (35.1)	45 (14.8)	153 (50.2)
I could easily discuss HIV-antibody testing with my family.	49 (16.1)	35 (11.5)	221 (72.5)
I consider going for HIV counselling and testing extremely frightening	74 (24.3)	44 (14.4)	187 (61.3)
Testing and counselling is a pleasant experience	87 (28.5)	45 (14.8)	173 (56.7)
There is a possibility that I have HIV and AIDS	166 (54.4)	52 (17.0)	87 (28.5)
I may have had sex with someone who was at risk for HIV and AIDS	181 (59.3)	29 (9.5)	95 (31.1)
I am at risk of HIV and AIDS	99 (32.5)	38 (12.5)	168 (55.1)

*Significant at $p < 0.05$, †Fisher's Exact test used**Table 3:** logistic regression on socio-demographic factors influencing HCT uptake (n=305).

Predictors	B (regression coefficient)	95% CI for aOR			P-value
		aOR	Lower	Upper	
Marital status					
Married (Reference category)	1				
Single	1.45	4.265	1.866	9.746	0.001*
Relationship	1.996	7.361	1.768	30.656	0.006*
Divorced	0.137	1.147	0.096	13.75	0.914
Academic class					
First year (Reference category)					
Second year	1	2.718	1.124	6.575	0.027*
Third year	-0.014	0.986	0.332	2.925	0.979
Midwifery	0.472	1.604	0.537	4.788	0.397
Age of participants					
19-24 years (Reference category)	1				
25-30 years	-1.265	0.282	0.14	0.567	0.001*
31-50 years	-0.493	0.611	0.171	2.184	0.448
Religion					
Islam (Reference category)	1				
Christian	1.662	5.272	1.839	15.112	0.002*
Constant	-2.537	0.079			0.262

*Statistical significance p value < 0.05 , aOR=adjusted Odds Ratio, CI=Confidence Interval**Table 4:** Facilitator items and factor loadings for the HIV Antibody Testing Scale (n=305).

Items	Factors				
	1	2	3	4	5
My friends would not treat me any differently if I were tested for HIV.	<i>0.787</i>	-0.03	0.122	0.041	-0.161
My friends would look down on me if I were tested for HIV	-0.768	-0.167	0.032	0.076	-0.008
My friends would not look down on me if I were tested for HIV	<i>0.653</i>	0.006	0.006	0.258	0.146
It would not bother me if someone I know sees me going to get an HIV test	<i>0.372</i>	-0.053	0.244	0.221	-0.278
My friends would support my decision to get an HIV test	<i>0.347</i>	-0.117	0.322	0.286	-0.038
I may have had sex with someone who was at risk for HIV and AIDS	-0.047	<i>0.82</i>	-0.058	-0.01	-0.079
There is a possibility that I have HIV and AIDS	0.065	<i>0.82</i>	-0.016	0.171	-0.054
I am at risk of HIV and AIDS	0.061	<i>0.742</i>	0.136	-0.069	0.065
HIV tests give accurate results	-0.003	0.123	<i>0.71</i>	-0.012	-0.128
I would be comfortable talking to an HIV counselor about personal behaviors that place me at risk for HIV infection	0.032	-0.074	<i>0.689</i>	0.097	0.245
It is extremely useful to test for HIV	0.075	0.088	<i>0.678</i>	0.063	0.228
I could easily discuss HIV-antibody testing with my family.	0.164	-0.089	<i>0.494</i>	0.385	-0.119
I trust the HIV test counselors and nurses to keep my information confidential.	0.232	0.044	-0.002	<i>0.746</i>	0.222
HIV test information is kept very confidential by the medical staff who do the testing.	0.068	0.068	0.089	<i>0.71</i>	0.079
Testing and counselling is a pleasant experience	-0.031	0.014	0.159	<i>0.621</i>	-0.326
I consider going for HIV counselling and testing extremely frightening	-0.006	-0.023	0.048	-0.108	<i>0.764</i>
I would like to be alone when doing the test	-0.081	-0.06	0.219	0.283	<i>0.61</i>

The factors were interpreted as relating to: 1 = support of friends/partners, 2 = positive perceived susceptibility, 3 = personal concerns about HCT, 4 = assured confidentiality and support and 5 = fear about HCT, disclosure, stigma & discrimination. Items are assigned to the scale with highest loading (in bold).

Table 5: Barrier items and factor loadings HCT uptake on HCT Scale (n=305).

Items	Factors				
	1	2	3	4	5
I would not get tested for HIV because I would be asked information that was too personal	0.722	0.135	0.075	0.09	0.14
I do not have time to get an HIV test	0.69	0.315	-0.22	-0.057	-0.085
I would be embarrassed if my friends found out that I had decided to have an HIV test.	0.675	0.099	0.328	0.206	0.108
I am afraid someone would find out I was tested for HIV	0.657	0.047	0.374	0.21	0.057
My friends would treat me badly if I were tested for HIV	0.417	-0.041	0.389	0.255	-0.117
Anyone who is tested for HIV is dirty.	0.077	0.759	0.068	0.039	0.043
Admitting that you should be tested for HIV means that you have engaged in immoral behavior	0.028	0.668	0.126	0.133	-0.054
HIV testing is unnecessary for me, as I feel healthy	0.183	0.579	0.058	-0.272	0.018
I consider going for HIV counselling and testing extremely humiliating	0.268	0.509	-0.072	0.342	0.231
People assume that everyone who is tested for HIV is infected with HIV.	-0.015	0.181	0.717	-0.105	0.092
People would assume I have HIV if I decided to get tested	0.4	-0.096	0.704	-0.016	-0.147
I am afraid that if I were tested for HIV, my name would go into public record	0.102	0.374	0.537	0.265	0.232
I would not want anyone to know if I got an HIV test.	0.221	-0.058	-0.088	0.784	0.036
It would be embarrassing to get tested for HIV	0.082	0.384	0.34	0.566	-0.051
HIV-antibody testing is not really confidential	0.088	0.027	0.044	0.009	0.948

Table 6: Socio-demographic factors associated with attitude towards HCT (n=305).

Variable	N	Mean(SD)	Statistic value	p
Age			0.411**	0.663
19-24 years	177	113.61 (±11.54)		
25-30 years	88	113.86 (±13.653)		
31-50 years	40	115.55 (±12.098)		
Gender			-0.33^	0.742
Female	186	113.75 (±11.728)		
Male	119	114.23 (±13.028)		
Family type			1.104^	0.271
Nuclear Family	157	114.69 (±13.153)		
Extended Family	148	113.14 (±11.162)		
Residence			0.885^	0.028*
Urban	174	115.27 (±12.034)		
Rural	131	112.17 (±12.315)		
Ethnicity			1.111**	0.355
Mandinka	125	113.24 (±12.498)		
Fula	68	114.49 (±11.475)		
Wolof	30	111.8 (±14.414)		
Jola	36	118.19 (±12.259)		
Manjago	14	116.5 (±12.126)		
Sarahuli	7	107.29 (±12.619)		
Serere	13	113.69 (±7.398)		
Aku	7	111.71 (±10.657)		
Foreigner	5	112 (±10.954)		
Marital Status			3.543**	0.015*
Married	88	114.86 (±12.381)		
Single	199	114.17 (±11.842)		
In a relationship	13	103.38 (±13.314)		
Divorced	5	115.8 (±13.312)		
Year of Study			0.307**	0.820

Discussion

In our study, majority (58.4%) of the students had been tested for HIV at least once. This is much higher than the 8 % for males and the 14 % for females reported as reproductive-aged adults (ages 15-49) in the 2018 Multiple Indicator Cluster Survey (MICS) in The Gambia [14]. In addition, the proportion of those who tested was higher than the 3% for males and the 9% for women reported for young adults (15-24 years) in the same national survey. The finding followed a similar trend among studies conducted in Ghana and also in Zambia [23,24] among nursing students. It also appears that even among would-be nursing professionals HCT uptake was not at 100%, which signifies that there are some factors that hinders HCT uptake as consistent with a study conducted among university students in Nigeria, only about half (50.7%) of the participants had HCT [25].

The finding also showed that majority of the students were aware of the importance of HCT in the prevention HIV/AIDS and yet still the proportion of those who tested was not impressive. In response to this, we acknowledge that when and where to test for HIV did not reflect expected behaviour of seeking for HCT services and this disparity has been reported in other studies [13] and also a study conducted in Zambia further showed that despite majority of students who demonstrated willingness to undergo HCT, few actually had the test unless there is an underlying illness [26]. All the nursing schools from where the participants are drawn from have very close proximity to health facilities that offer HIV counseling and testing services yet still there was no hundred percent HCT utilization. The explanation to this scenario could be as a result of self-perception of not been susceptible to the disease or that feeling of absolute well-being which is common among young adults.

The result showed that age, gender, marital status, religion, academic year, school of attendance and residence (LGA) were strongly associated with HIV uptake. Our observations seemed to show that HCT utilization is better in older age at School since the age group between 25-30 years and 31 years score better than between 19-24 age group. However, this in contrast to sub-Saharan Africa that found younger adults more willing than

3rd Year Class	42	113.88 (±15.658)		
2nd Year Class	110	113.09 (±11.909)		
1st Year Class	88	114.58 (±8.907)		
Midwifery Class	65	114.54 (±14.217)		
School			0.790**	0.455
RN School	133	112.94 (±12.82)		
CHN School	97	114.62 (±10.919)		
SEN School	75	114.83 (±12.793)		
Parent alive			0.355**	0.786
One alive	108	114.51 (±12.344)		
Yes	154	113.36 (±12.558)		
Both alive	20	115.8 (±9.266)		
Divorced	23	113.48 (±12.176)		
Education father			1.380**	0.249
Tertiary	97	113.3 (±13.718)		
None	145	113.73 (±11.689)		
Primary	30	112.63 (±9.357)		
Secondary	33	117.91 (±11.977)		
Education Mother			0.502**	0.681
Tertiary	37	112.59 (±15.179)		
None	176	113.78 (±12.19)		
Primary	49	113.86 (±11.079)		
Secondary	43	115.84 (±10.987)		
Wealth index			1.225**	0.300
Middle	148	113.35 (±11.795)		
Lowest	67	112.19 (±12.828)		
Second	77	116.23 (±11.343)		
Fourth	7	115 (±12.138)		
Highest	6	117.17 (±23.6)		
LGA			0.431**	0.882
Brikama	107	113.81 (±12.709)		
Kanifing	91	114.43 (±11.611)		
Kerewan	15	113.67 (±14.171)		
Basse	22	110.05 (±15.126)		
Janjanbureh	22	115.82 (±12.006)		
Banjul	18	115.06 (±7.008)		
Mansakonko	24	113.79 (±13.012)		
Kuntaur	6	114 (±8.944)		

older people to be tested for HIV [27,28]. These could be due to the fact that as respondents grow in age, they engage into conjugal relationships that may increase their perceived susceptibility and vulnerability to HIV.

Our finding showed more male students testing for HIV compared to females. These results were the opposite or not in alignment with the national trend where males are less likely than females to have been tested for HIV in The Gambia and in other studies from across sub-Saharan Africa [29]. Moreover, this finding is non-congruent with studies done in Ghana and Kenya [30,12]. The utilization of HCT services among females as realized in most sentinel surveys or demographic health surveys could be due to the ongoing prevention of mother-to-child transmission program during antenatal services in The Gambia that requires every pregnant female who reports to the health facilities to undergo HIV testing as a protocol of the policy. Furthermore, gender inequity poses a great challenge to HCT up-

take especially in sub-Saharan nations (Ghana, Gambia, Nigeria, Uganda etc.), where the sole authority or power and responsibility of healthcare-seeking behaviours depend on men [31]. In the Gambia men may refuse HCT service as a show of strength or dominance or self-confidence which is very typical in many African homes. If the wife suggests HCT, this may be seen as undermining the roles of the man as a decision maker.

Furthermore, student nurses in the third year of study and midwifery class were two times more likely to utilize HCT services when compared to those in year 1 and 2. These findings suggest that the higher the educational level of student, the more likely that they will utilize HCT services. This finding is similar to a study done in Addis Ababa which predicts that educational year is positively correlated with HCT uptake [32]. This finding could be interpreted as the more time students spent on the educational ladder the more likely they will utilize HCT services. In addition, our study showed that marital status was significantly associated with HCT uptake. Our finding also showed that divorcees were equally more likely to undergo HIV counselling and testing than students who are single. This finding corroborated with other studies [33] across Africa among student nurses. In many parts of sub-Saharan Africa, the fear of divorce or broken marital relationships which may result to potential abandonment or even violence can serve as a potential deterrent to the uptake of HCT services [34]. Our findings also aligned with findings from a Jamaican study which revealed that married persons were more likely to report previous HIV testing [35] than unmarried persons. The high figures among married and divorcees may be partially due to the parent to mother-child transmission program which offers routine HCT to services to all pregnant mothers and partners. Yet still it could be related to the fact that young unmarried persons see themselves as less susceptible to HIV.

On knowledge on HCT, the study findings revealed that students who are aware were more likely to utilize HCT services compared to those who are not. This finding was similar to studies done in Nigeria and Uganda among youths [36,37]. This finding further emphasized the need to strengthen awareness campaigns about HCT services rather than concentrate solely on facts about HIV/AIDS. With regards to preferred person for HCT, most of the participants preferred trained counselor to doctors and nurses. This finding was different from other similar studies in Ethiopia [38]. Our study also showed majority of participants willing to go for HCT in the future. This finding was similar to a study conducted in Mersa Town of Harbu district, Ethiopia [39]. This high proportion of students willing to undergo HCT was not parallel with the actual use of HCT services.

On attitude towards HCT, our findings showed a little over half of the student nurses and midwives had positive attitude towards HCT uptake. The proportion of students with positive attitude towards HCT was far lower compared to a study conducted in North West Ethiopia which revealed 73.3% of student having positive attitude towards HCT uptake [39]. This disparity between the current study and other studies with high proportion of positive attitude towards HCT could be as a result of numerous factors. The explanation for the unimpressive proportion in this current study could be related to inadequate knowledge on HCT as demonstrated in our study. Our study also revealed that students with stigmatizing attitudes were less likely to utilize HCT than those who had positive attitude towards HCT [40]. This is because stigma is seen as a barrier to HCT uptake which makes people less enthusiastic to seek for HCT. The finding aligns

with finding from a study done in Mersa Town, Ethiopia among adults which indicated that persons with stigmatizing attitude were less likely to utilize HCT services compared to those with positive attitude [39]. Similarly, a study in South Africa also reported consistent finding [15].

On the socio-demographic variables associated with attitude towards HCT uptake, previous residence and marital status were significantly associated with attitude towards HCT. Finding showed that students who reside in rural areas showed more stigmatizing attitudes towards HCT compared to those from urban areas. Also, result showed that married students showed more positive attitude towards HCT compared to students who are single or in a relationship. This finding was similarly with a study conducted among students in Addis Ababa University [41]. The plausible explanation for this observation could be related to the fact that students from rural areas lack good information about HCT due to adequate access news outlets/media. Also, married students are matured and may have undergone HCT before.

Aside from the socio-demographic factors explored in this present study, we also sought to determine other influencing factors associated HIV uptake. Results from factor analysis showed that HCT uptake was associated with five main barriers. These barriers are ranked as follows: concerns about negative stereotyping from friends, concerns of people, personal concerns about HCT, privacy & attitude of health staff and lack of confidentiality and support. These findings were similar to a study conducted in Kenya by Rose Mwangi et al., 2012 [18]. In this study the most important barrier associated with HIV uptake was personal concerns and concerns of friends which is strongly associated with stigma and discrimination. Therefore, stigma and discrimination is a major barrier to HCT uptake among students. Therefore, when students have in mind that they will be stigmatized and discriminated for having a HIV test, there is a strong likelihood that they will stay away from undertaking HIV services. The second barrier is related to lack of adequate knowledge on the benefits of HCT. Hence participants see HCT as valueless and not worth undertaking, thus, it is crucial that youths are well educated on the importance and value of HCT so as to increase uptake. This was evident among those participants who did not undertake HCT in our current study who mentioned that self and partner mistrust were the main reason for not undertaking the HIV test. The third barrier observed in this study was peoples' concerns in which the major concern expressed was the risk of their status being exposed to sexual partners or other people within the community. These types of concerns are common throughout Africa [42,43]. Many studies showed that adults willing to be tested usually prefer to be counseled and tested by someone who does not know them hoping that their result will not be exposed [44]. Similarly, fear of rejection from partners or family members or friends is related to this particular barrier and very common in many setting in Africa including the Gambia. It is therefore imperative that testing is done in very secure health institutions where services are sought. The fourth factor or barrier is about privacy and attitude of staff at health facility which can hinder the uptake of HCT. The barrier is also reported in a study conducted in Ghana among nurses [45]. Health staff are supposed to show positive attitude towards clients and equally motivate and encourage them to seek for services to enhance good health. Negative attitude towards patients/clients could hinder the uptake of services. Similarly, maintaining privacy during service delivery is paramount and can encourage more clients to seek for services.

Therefore, lack of privacy and negative attitude of staff can seriously hinder the uptake of HCT services across the different cohorts of population especially the youths. Lack of confidentiality is another barrier reported in this current study. This barrier was also mentioned in a study conducted in Ghana among student nurses [45]. Lack of confidentiality can adversely decrease the uptake of HCT among trainee nurses and midwives. Clients generally feel insecure or worried that their private information may be shared with loved ones or exposed to the general public. This is very common especially in African country where secure data storage is still a challenge. This is also coupled with the fact that those offering the service may be known to you or your family members or friends hence the fear that the vital information may be shared with them.

On the facilitators associated with HIV counseling and testing, our study revealed that support from friends/partners or family members can greatly increase the confidence of persons opting to undergo HIV counseling and testing. This was one of the main facilitators been expressed by the students and it is paramount that more health education campaigns are focus on the youths to change their misconceptions about HCT so as to offer more support or motivation to those willing to undergo HCT. In many studies, individuals fear to undergo HCT because of lack of support from partners or friends who may taunt or stigmatize their action [46]. Health care seeking behaviour is strengthened or influenced by close partners especially in the Africa context especially among students in particular. Students tend to follow the foot step of their friends or family members in health related activities, hence their support or encourage is paramount. The second facilitator that can motivate people to undergo HCT was having that positive feeling of been susceptible to HIV infection. Based on the Health Belief Model, people tend to change a behavior if they felt susceptible to a disease, therefore people must be knowledgeable on the modes of HIV transmission and be able to assess their actions and take the appropriate measures to change positively. Many studies revealed that people who feel less susceptible do not undergo HIV testing [47]. Therefore, it is crucial that we discourage individuals from that false belief of low susceptibility or the belief that 'they are healthy' and does not need a HIV test. The best way to assess your behavior in terms of HIV infection is to go for HCT therein you will be assisted by a trained counselor to assess your actions. As young adults who are active and with strong immune system coupled with little or good knowledge on HIV/AIDS, students would tend to believe that they do not exhibit the signs of HIV hence there is no need for testing. Many may regard testing as a waste of time bearing in mind that there is no cure for a positive test. Our study also revealed that HCT uptake can be increase if personal and other people's concerns are addressed. The finding showed that when an individual value the important of HCT, there is a likelihood that they will undergo HIV testing. The finding is supported by the health belief model which states that if individuals appreciate the benefits of a particular behaviour they tend to adopt interventions recommended to bring about the change. Therefore, individual concerns with regards to HCT in a positive manner can promote or influence HCT uptake among student nurses and midwives. Many of the students are far away from family home and are considered mature enough to make their own decision hence family influence may not play a major role. This finding was also supported by results from a study conducted among university students in four African countries which showed that going for HCT was related to: general concerns, trust and support, and

fears. Additionally, in the same study, 'Friends concerns' were also associated with the intention to go for an HIV test [48].

On the fourth facilitator, which is related to assured confidentiality, support and privacy are paramount in the utilization of HCT services among student nurses. This is an isomer to the results of a study conducted in Ghana where the importance of privacy and confidentiality in HCT uptake was evident [45]. It demonstrated that clients and potential users of services are uncomfortable with the quality of care given by some health workers, especially as they overtly and covertly breached confidentiality about their clients' health status. This has compelled many patients and potential users of the services to adopt a modus vivendi that provides them access to some care services while protecting their identity. Therefore maintaining confidentiality is key to increase in uptake of HCT among student nurses and other cohorts.

The fifth facilitator was on privacy and fear in knowing their HIV status. Therefore, adequate privacy should be accorded to clients during services delivery. The mere presence of a person at the HIV counselling centre or clinic is enough for the person to be labelled as or suspected to be HIV patient. It demonstrates that stigmatization may occur not only in the community but also overtly or covertly, in the health facility itself.

Practical implications of The Study

The role or relevance of nursing sector of the health force is crucial in any effective HIV/AIDS programs. Thus, nursing students should be regarded as key partners in the fight against HIV and AIDS. We recommended that all key players in the fight against HIV/AIDS should be encouraged to regularly undergo HCT, so they will be confident enough to give advice to others to seek for HCT services. Moreover, there is a great need to embark on extensive HIV/AIDS educational programs so that nurses can be agents of change. Despite the limitations, the study provides useful information on education and strategic planning for future HIV programs. The study also provide an insight on the prevalence of HIV counseling and testing uptake among nursing students and midwives, which is crucial for future strategic planning.

Limitations of this Study

In our study, we were faced with some challenges that ought to be taken into considerations. As for such, the closed-ended responses of the self-administered questionnaire did not give opportunity for respondents to express their opinions about other influencing factors associated with HCT. In addition, information provided by the respondents may be bias due to misunderstanding of questions or as a result of giving false information.

Conclusion

From our study, socio-demographic factors such as sex, age, year in study, wealth index and to some extent marital status are found strongly correlating with the uptake of HCT service among our study participants. In addition to this, the uptake of HCT service are hindered by facilitators and multifactorial barriers. Moreover, we found that decrimination, stigma, friends and family contributing stereotypes, privacy, which might arise from personal concerns, perceived susceptibility and confidentiality are the key barriers to HCT uptake. It is suprising that despite a good number of students do attend HCT, still the utilization rate was not impressive as expected from would-be nurses. Simi-

larly, knowledge on HCT was not significantly high as expected. However, students showed positive attitude towards HCT.

From our findings, we recommend that health education programs specific on the benefits of HCT and other relevant topics that are being undertaken by different stakeholders in the fight against HIV/AIDS through drama and entertainment should be strengthened. Also, HIV/AIDS educational programs should be incorporated into school curricula so as to increase knowledge on HCT among students and clear misconceptions regarding HIV/AIDS and HCT. Furthermore, HCT services should be provided in nursing schools to improve access and thus limit stigma and discrimination that may prevail in some health institutions.

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and material: Not applicable

Competing interests: The authors declare no competing interest

Funding: No funding is associated with this project

Authors' contributions: SM and Prof Sun Mei conducted the study, OB, SM and Prof Sun Mei designed the study, SM and AB analyzed the data and all the authors proofread the paper.

References

1. Kwabong GD, Boateng D, Agyei-Baffour P, Addy EA. Health service barriers to HIV testing and counseling among pregnant women attending Antenatal Clinic; A cross-sectional study. *BMC Health Serv Res.* 2014; 14.
2. Deblonde J, De Koker P, Hamers FF, Fontaine J, Luchters S, et al. Barriers to HIV testing in Europe: A systematic review. *Eur J Public Health.* 2010; 20: 422-432.
3. Peltzer K, Matseke G, Mzolo T, Majaja M. Determinants of knowledge of HIV status in South Africa: Results from a population-based HIV survey. *BMC Public Health.* 2009; 9: 1-11.
4. Reis C, Heisler M, Amowitz LL, Moreland RS, Mafeni JO, et al. Discriminatory attitudes and practices by health workers toward patients with HIV/AIDS in Nigeria. *PLoS Med.* 2005; 2: 0743-0752.
5. Kalyanshetti S, Nikam K. A study of knowledge of HIV/AIDS among nursing students. *Int J Med Sci Public Heal.* 2016; 5: 1209.
6. Charles MP, Kweka EJ, Mahande AM, Barongo LR, Shekalaghe S, et al. Evaluation of uptake and attitude to voluntary counseling and testing among health care professional students in Kilimanjaro region, Tanzania. *BMC Public Health.* 2009; 9: 1-9.
7. Patton GC, Coffey C, Sawyer SM, Viner RM, Haller DM, et al. Global patterns of mortality in young people: a systematic analysis of population health data. *Lancet.* 2009; 374: 881-892.
8. Statistics TGB of. The Gambia Demography and Heath Survey. 2014; 2014-2015.
9. The impact of Voluntary.
10. SADC. Assessment Report on the Status of HIV Testing and Counselling Policies in the SADC Region. 2009.
11. Journal SA, Health P, Rehanilitation M, Student MPH. Accept-

- ability and Barriers To Uptake of Hiv Testing and Counseling Among Students of Tertiary Institutions in Owo Ondo State Nigeria. 2014; 1: 1-17.
12. Paul N, Muewa M. Public Health in the School of Public Health , Kenyatta University A Research Thesis submitted in partial fulfillment of the requirement for the award of the Degree of Master of. 2014.
 13. Obermeyer CM, Osborn M. The utilization of testing and counseling for HIV: A review of the social and behavioral evidence. *Am J Public Health.* 2007; 97: 1762-1774.
 14. Gambia. Central Statistics Dept., UNICEF. MICS, The Gambia. 2018:96, xxv p.
 15. Kalichman SC, Simbayi LC. HIV testing attitudes, AIDS stigma, and voluntary HIV counselling and testing in a black township in Cape Town, South Africa. *Sex Transm Infect.* 2003; 79: 442-447.
 16. Daniyam CA, Agaba PA, Agaba EI. Original Article Acceptability of voluntary counselling and testing among medical students in Jos. Nigeria. *J Infect Dev Ctries.* 2010; 4: 357-361.
 17. WHO. Towards the universal Access: Scaling up priority HIV/AIDS interventions in the health sector. *Unaided Who, Unicef.* 2009.
 18. Mwangi RW, Ngure P, Thiga M, Ngure J. Factors influencing the utilization of Voluntary Counselling and Testing services among university students in Kenya. *Glob J Health Sci.* 2014; 6: 84-93.
 19. Zade Beg No Title Basics of food chemistry by Jan Daman. 2014.
 20. Boshamer CB. BKA scale to measure attitudes about H testing: development and psychometric validation. *AEP* 1999 O-13. P 10555624. A scale to measure attitudes about HIV-antibody testing: development and psychometric validation. *J Clean Prod.* 2015; 86: 311-322.
 21. Sambah F, Elvis J, Jr H, Hormenu T, Ahinkorah BO, et al. Determinants of HIV Testing and Counseling Utilization among Trainee Nurses and Midwives in Central Region of Ghana. 2019; 2: 1-10.
 22. Gaskin CJ, Happell B. On exploratory factor analysis: A review of recent evidence, an assessment of current practice, and recommendations for future use. *Int J Nurs Stud.* 2014; 51: 511-521.
 23. Bhoobun S, Jetty A, Koroma MA, Kamara MJ, Kabia M, et al. Facilitators and barriers related to voluntary counseling and testing for HIV among young adults in Bo, sierra leone. *J Community Health.* 2014; 39: 514-520.
 24. Child Y, Practices F, The IN, End E, Freetown OF, Leone S. Sierra Leone Journal of Biomedical Research. 2016; 8: 4-11.
 25. Daniyam CA, Agaba PA, Agaba E. Acceptability of voluntary counselling and testing among medical students in Jos, Nigeria. *J Infect Dev Ctries.* 2010; 4: 357-361.
 26. Jürgensen M, Tuba M, Fylkesnes K, Blystad A. The burden of knowing : balancing benefits and barriers in HIV testing decisions . a qualitative study from Zambia. 2012; 10-13.
 27. Negin J, Nemser B, Cumming R, Lelera E, Ben Y, Paul A. HIV Attitudes , Awareness and Testing Among Older Adults in Africa. 2012; 63-68.
 28. Tabana H, Doherty T, Swanevelde S, Lombard C, Jackson D, et al. Knowledge of HIV status prior to a community HIV counseling and testing intervention in a rural district of south Africa : results of a community based survey. *BMC Infect Dis.* 2012; 12: 73.
 29. Magadi MA. Understanding the gender disparity in HIV infection across countries in sub-Saharan Africa : evidence from the Demographic and Health Surveys. 2011; 33: 522-539.
 30. Oppong Asante K. HIV/AIDS knowledge and uptake of HIV counselling and testing among undergraduate private university students in Accra, Ghana. *Reprod Health.* 2013; 10: 1-8.
 31. Grant E, Logie D, Masura M, Gorman D, Murray SA. AIDS Care : Psychological and Socio-medical Aspects of AIDS / HIV Factors facilitating and challenging access and adherence to antiretroviral therapy in a township in the Zambian Copperbelt : a qualitative study. 2008; 37-41.
 32. Addis Z, Yalew A, Shiferaw Y, Alemu A, Birhan W, et al. Knowledge , attitude and practice towards voluntary counseling and testing among university students in North West Ethiopia : a cross sectional study. 2013; 1-8.
 33. Dirar A, Mengiste B, Kedir H, Godana W. Factors contributing to voluntary counselling and testing uptake among youth in colleges of Harar , Ethiopia. 2013; 1: 91-96.
 34. Mlay R, Lugina H, D SBP. AIDS Care : Psychological and Socio-medical Aspects of AIDS / HIV Couple counselling and testing for HIV at antenatal clinics : Views from men , women and counselors. 2014; 37-41.
 35. Norman LR. ORIGINAL RESEARCH HIV testing practices in Jamaica. 2006; 231-242.
 36. Babalola S. Readiness for HIV Testing among Young People in Northern Nigeria : The Roles of Social Norm and Perceived Stigma. 2007; 759-769.
 37. Gage AJ, Ali D. AIDS Care : Psychological and Socio- medical Aspects of AIDS / HIV Factors associated with self-reported HIV testing among men in Uganda. 2010; 37-41.
 38. Faculty Of Medicine Department Of Community Health Determinants Hiv-Vct Acceptance In Gondar Town , Northwest Ethiopia : A Case Control Study . By : Anwar Yibrie Endris , Md Thesis Submitted To The Of Parital Fulfillment Of The Requirments For The Degree.
 39. Tsegay G, Edris M, Meseret S. Assessment of voluntary counseling and testing service utilization and associated factors among Debre Markos University Students , North West Ethiopia : a cross-sectional survey in 2011. 2013.
 40. Access O. Patterns of sexual risk behavior among undergraduate university students in Ethiopia: a cross-sectional study. 2012; 8688: 1-9.
 41. Regassa N. KSA and practices on H preventions among students of higher education institutions in E the case of AAUEAJPH 2011; 8: 141-154. TAttitudes and practices on HIV preventions among students of higher education institutions in Ethiopia: the case of Addis Ababa University. *Drugs.* 2017; 77: 1295-1311.
 42. Njau B, Covin C, Lisasi E, Damian D, Mushi D, et al. A systematic review of qualitative evidence on factors enabling and deterring uptake of HIV self-testing in Africa. *BMC Public Health.* 2019; 19.
 43. Fylkesnes K, Siziya S. A randomized trial on acceptability of voluntary HIV counselling and testing. *Trop Med Int Health.* 2004; 9: 566-572.
 44. Angotti N, Bula A, Gaydosh L, Zeev E, Thornton RL, et al. Social Science & Medicine Increasing the acceptability of HIV counseling and testing with three C ' s : Convenience , confidentiality and credibility q. *Soc Sci Med.* 2009; 68: 2263-2270.
 45. Dapaah JM, Senah KA. HIV/AIDS clients, privacy and confidentiality; The case of two health centres in the Ashanti Region of Ghana. *BMC Med Ethics.* 2016; 17: 1-10.
 46. Boshamer, C. B., & Bruce KE. A scale to measure attitudes about H, (5) testing: D and psychometric validation. *AEPO* 11, 400-

413. A scale to measure attitudes about HIV-antibody testing: development and psychometric validation. *Int J Adv Couns.* 2002; 24: 193-203.
47. Denison JA, Reilly AE, Schmid GP, Kennedy AE, Sweat MD. HIV Voluntary Counseling and Testing and Behavioral Risk Reduction in Developing Countries : A Meta-analysis, 1990-2005. 2008; 363-373.
48. Meiberg AE, Bos AER, Onya HE, Schaalma HP. FEAR OF STIGMATIZATION AS BARRIER TO VOLUNTARY HIV COUNSELLING AND TESTING IN SOUTH AFRICA Annemarie E. Meiberg 1 , Arjan E. R. Bos 2 , Hans E. Onya 3 , Herman P. Schaalma 1. 2008; 5: 49-54.