



# Molecular Detection of Aflatoxin Producing *Aspergillus* Species Isolates in Some Chicken Meat Cuts in Gharbiya Governorate, Egypt

Shaltout FA<sup>1</sup>; Heikal GI<sup>2</sup>; Ghanem AM<sup>3\*</sup>

<sup>1</sup>Food Hygiene and Control Department (Meat Hygiene), Faculty of Veterinary Medicine, Benha University, Egypt.

<sup>2</sup>Food Hygiene Department, Animal Health Research Institute (Tanta branch), ARC, Egypt.

<sup>3</sup>Food Security Sector, Egyptian Armed Forces, Egypt.

\*Corresponding Author(s): **Ghanem AM**

Food Security Sector, Egyptian Armed Forces, Egypt.

Email: fahimshaltout@hotmail.com

## Abstract

Contamination with fungi and their toxins is considered one of the most dangerous hidden pollutants that threaten the health of the consumer. The presence of mycotoxins in various foods has been recorded, despite their apparent safety for human consumption. Therefore, the current study was conducted to evaluate the prevalence of *Aspergillus* species by culture method; and aflatoxin-producing genes molecularly in total of 75 random samples of chicken cuts represented by wing, breast and thigh (25 of each) that were collected from various groceries and poultries shops located at Gharbiya governorate, Egypt. Results of culture and isolation techniques revealed detection of *Aspergillus* sp. in 36, 48 and 44% of the examined wing, breast and thigh samples, respectively. Moreover, microbiological identification of the isolated strains showed presence of *A. niger*, *A. flavus*, *A. fumigatus*, *A. terreus* and *A. parasiticus* in 16, 13.3, 10.6, 1.3 and 1.3% of the total population of the examined samples. Molecular detection of some aflatoxin production regulating genes (*OmtA*, *Nor1* and *Ver1*) in ten *Aspergillus* sp. isolates revealed their detection in 8/10 (80%), 8/10 (80%) and 7/10 (70%) represented by positive bands at molecular weight of 1024, 400 and 537 bp, respectively. Referring to the recorded results, chicken cuts may possess a great silent hazard to the human being under improper good manufacturing practices and inadequate hygienic conditions during handling and storage.

Received: Dec 08, 2021

Accepted: Feb 15, 2022

Published Online: Feb 18, 2022

Journal: Annals of Epidemiology and Public health

Publisher: MedDocs Publishers LLC

Online edition: <http://meddocsonline.org/>

Copyright: © Ghanem AM (2022). This Article is distributed under the terms of Creative Commons Attribution 4.0 International License

**Keywords:** *Aspergillus* species; Chicken meat cuts; cPCR; Egypt.



## Introduction

Chicken meat and meat products production in developing countries plays an essential role in supporting food security and poultry meat demands [1].

Contamination of meat products with molds can occur during different preparation stages during slaughtering under bad hygienic conditions using contaminated water or by adding contaminated spices with mold spores or during packing, handling, transportation and storage [2].

Contamination of meat with *Aspergillus* species, especially Flavi section, is one of the most hazardous microbial contamination as the majority of *Aspergillus* species are able to aflatoxins production that can cause diseases associated with aflatoxin poisoning and carcinogenic effects [3].

Acute aflatoxin poisoning may lead to death as was recorded in Kenya in 2004 [4], while chronic poisoning may lead to various recorded mutagens and cancers [5].

*Aspergillus* sp. was classified into two groups depending on their toxigenic impacts on food and human health; 1st group includes the aflatoxigenic species such as *A. flavus* and *A. parasiticus*, while the 2nd group contains the non-aflatoxin-producing species such as *A. tamarisii* and *A. oryzae* [6].

Molecular analyses have been used to confirm aflatoxin productivity of *Aspergillus* species isolates. *omtA*, *nor1* and *ver1* genes are from the commonly used genes encoded *aflP*, *aflD* and *aflM* toxins detection in food items [7] yield an accurate, rapid and reliable records of toxigenic *aspergillus* species especially in food chain [8].

Therefore, the main target of the current study was to investigate the presence of toxigenic *aspergillus* species in some chicken meat cuts collected from Gharbiya Governorate markets, Egypt.

## Material and methods

### Collection of samples

A total of seventy-five random samples of raw chilled chicken wing, chicken thigh, chicken breast (25 of each) were collected from different local poultries shops and different supermarkets at Gharbia governorate, Egypt. Samples were taken aseptically in polyethylene bags and were transferred to the laboratory in ice box for mycological examination.

### Preparation of samples [9]

Twenty-five grams from each sample were carefully and aseptically homogenized in blender after mixing with 225 ml of sterile peptone water 0.1% to form a dilution of 1:10, from which tenth fold serial dilutions were prepared.

### Determination of aspergillus species

**Culture of the prepared samples:** Was performed according to ISO [10], where 0.1ml of the previously prepared serial dilutions was spread by mean of sterile L-shape glass rod over two Petri-dishes contained solidified Dichloran Rose Bengal agar with Chloramphenicol (DRBC) then were incubated at upright position at 25°C for 5 -7 days.

**Identification of isolated strains:** Was performed according to Pitt and Hocking [11] macroscopically and microscopically as recorded in (Table 1).

**Molecular detection of some aflatoxin producing genes of some isolated *Aspergillus* strains by cPCR**

### Oligonucleotide primers used in cPCR

Three pairs of *omtA*, *nor1* and *ver1* primers were prepared and collected from Metabion (Germany). Their special sequence and amplify certain products as were be displayed in (Table 2).

**Mycological DNA:** Was extracted following QIAamp DNeasy Plant Mini kit Catalogue no. 69104.

**Preparation of master mix and thermal profile:** Was adapted according to the manufacturer instructions (Emerald Amp GT PCR mastermix (Takara) Code No. RR310A).

**Table 1:** Morphological character of *Aspergillus* species [11].

	Colony Diameter (mm)	Texture	Surface color	Reverse color	Stipes	Vesicles	Seriation	Conidia	Colonial head/ cleistotheca
<i>A. flavus</i>	65-70	Floccose powdery or granular	Greenish yellow	Pale brown	Rough hyaline	Globose or subglobose	Biseriate	Globose to ellipsoid	Radiating head
<i>A. fumigatus</i>	40-70	Velvety to powder	Blue with white margin	Slight green	Smooth hyaline	Clavate	Uniseriate	Globose or subglobose	Columnar head
<i>A. nidulans</i>	50-65	Velvety	Green	Brown	Smooth brown	Pyriform	Biseriate	Globose rough	Radiate Head/hulla Cells ascis
<i>A. niger</i>	50-70	Granular of powdery	Black	Pale yellow	Smooth Yellow to brown	Round	Biseriate	Globose brown	Round head
<i>A. terreus</i>	40-60	Powdery	Sandy to brown	Pale brown	Smooth hyaline	Round to pyriform	Biseriate	Globose to ellipsoid	Columnar head

**Table 2:** Oligonucleotide primers sequences.

Gene	Sequence (5'-----3')	Amplified product (bp)	Reference
<i>omtA</i>	GGCCCGGTTCTTGCTCCTAAGC	1024	[12]
	CGCCCAGTGAGACCCTTCCTCG		
<i>nor1</i>	ACCGTACTACGCCGCACTCTCGGCAC	400	
	GTTGGCCGCCAGCTTCGACACTCCG		
<i>ver1</i>	GCCGCAGGCCGCGGAGAAAGTGGT	537	
	GGGGATATACTCCGCGACACAGCC		

**Results**

As recorded in (Table 3). *Aspergillus* sp. was detected in 32 (42.6%) of the total examined samples. In detail, breast samples recorded the highest contamination level (48%); followed by thigh and wing samples, respectively.

**Table 3:** Prevalence of *Aspergillus* species in the examined chicken meat cut samples (n= 25 of each).

Samples	Wing		Breast		Thigh		Total	
	No.	%*	No.	%*	No.	%*	No.	%**
<i>Aspergillus</i> sp.	9	36	12	48	11	44	32	42.6

\*%: Prevalence in Relation to the Number of Each Sample (25).

\*\*%: Prevalence in Relation to the total Number of Samples Population (75).

Regarding with the genus identification, *A. niger* had the highest detection levels (16%) in the examined samples (Table 4).


**Table 3:** Prevalence of identified aspergillus sp. in the examined chicken meat cuts (n= 25 of each).

Samples	wings		breast		thigh		Total	
	No.	%	No.	%	No.	%	No.	%
<i>Aspergillus</i> spp.								
<i>A. niger</i>	4	16	6	24	2	8	12	16
<i>A. flavus</i>	2	8	4	16	4	16	10	13.3
<i>A. fumigatus</i>	2	8	1	4	5	20	8	10.6
<i>A. terreus</i>	1	4	-	-	-	-	1	1.3
<i>A. parasiticus</i>	-	-	1	4	-	-	1	1.3


**Table 4:** Prevalence of aflatoxin producing genes in *A. flavus* isolates from the examined samples (n= 10).

Sample	<i>omtA</i>	<i>nor1</i>	<i>ver1</i>
1	+	+	+
2	-	-	-
3	+	+	+
4	+	+	+
5	+	+	+
6	+	+	+
7	+	+	-
8	-	-	-
9	+	+	+
10	+	+	+


Referring to the obtained results of molecular detection of some aflatoxin producing genes as recorded in (Table 5). and (Figure 1 to 3); *omtA*, *nor1* and *ver1* genes were detected in 8/10 (80%), 8/10 (80%) and 7/10 (70%) of the examined *A. flavus* isolates, respectively. presence of these genes indicated the Producibility of the examined strain for aflatoxins P, D and M, respectively.



**Figure 1:** Agarose gel electrophoresis of cPCR of *omtA* (1024 bp) gene of *A. flavus*. Lane L: 100 bp ladder as molecular size DNA marker; Lane P.: Control positive *A. flavus* for *omtA* gene; Lane N.: Control negative; Lanes 1, 3, 4, 5, 6, 7, 9 and 10: Positive *A. flavus* for *omtA* gene; Lanes 2 and 8: Negative *A. flavus* for *omtA* gene.



**Figure 2:** Agarose gel electrophoresis of cPCR of *nor1* (400 bp) gene of *A. flavus*. Lane L: 100 bp ladder as molecular size DNA marker; Lane P.: Control positive *A. flavus* for *nor1* gene; Lane N.: Control negative; Lanes 1, 3, 4, 5, 6, 7, 9 and 10: Positive *A. flavus* for *nor1* gene; Lanes 2 and 8: Negative *A. flavus* for *nor1* gene.



**Figure 3:** Agarose gel electrophoresis of cPCR of *ver1* (537 bp) gene of *A. flavus*. Lane L: 100 bp ladder as molecular size DNA marker; Lane P.: Control positive *A. flavus* for *ver1* gene; Lane N.: Control negative; Lanes 1, 3, 4, 5, 6, 9 and 10: Positive *A. flavus* for *ver1* gene; Lanes 2, 7 and 8: Negative *A. flavus* for *ver1* gene.

## Discussion

Chicken meat and meat products comply an important source of human protein supplement all over the world because they provide good source of digestible protein, low cholesterol fat, essential amino acids, minerals, and different types of vitamins and minerals.

In Egypt, as well as human population increasing, demand of animal proteins also is increasing represents a serious challenge in which poultry industry plays an essential role in filling nutrition gap as a rapid and more economic source of proteins [13].

Mold contamination of meat and meat products have been considered a serious source of food spoilage resulting in different organoleptic changes in flavor, color, texture, odor referred mainly to the fungal deterioration especially in poor developing countries due to lack of hygienic measures during processing and handling [14].

Presence of mold in foods may be referred to the rapid, easy disperse and wide spread of the fungal spores which are abundant in the environment introducing food chain through dust, water, workers and equipment. Their presence in food samples is a serious public health concern, as these fungi may be associated with the production of mycotoxins [15].

Aspergillus species represents an important mycotic infection in public health concern as a human pathogen and as toxin-producing food contaminant. It releases a lot of spores which found in air, water, soil, plant debris, manure and animal feed. As fungal spore's growing, it secretes digestive enzyme and mycotoxins leading to food spoilage and human mycotoxicosis [16].

Referring to the recorded results in (Table 3), *Aspergillus* sp. was prominently detected in breast samples other than wings and thighs samples, which came in agree with the previously recorded results of Darwish et al., [17].

Shaltout et al., [18] who found that the examined breast samples were more contaminated with fungal infection than wing and thigh samples. While the current prevalence of aspergillus species in the examined samples came lower than those recorded by Hassan [19] who found *Aspergillus* sp. in all the examined samples (100%) collected from Gharbiya governorate, Egypt. Moreover, Abuzaid et al., [20] also detected *A. flavus* and *A. niger* in 40 and 80% of the examined sausage samples of chicken origin, respectively.

Referring to the obtained results of the microbiological identification of *Aspergillus* sp. isolates as recorded in (Table 4), they came in agree with the previously reported results by Darwish et al., [17] who found that *A. niger* was the predominant detected strain, followed by *A. flavus* and *A. parasiticus* in the examined samples of chicken cuts collected from Zagazig city, Egypt.

Some mold species can cause respiratory infections representing a significant risk for individual with severely weakened immune system [21]. Presence of mold in high incidence indicate bad hygienic measures adopted during handling, preparation and processing [22].

Mycotoxins are have been defined as naturally occurring secondary fungal metabolites produced in meat and meat products by direct growth of toxigenic molds such as *Aspergillus* species which produce Aflatoxins and Ochratoxins which threat public health due to their carcinogenic, hepatotoxic, nephrotoxic, teratogenic and mutagenic effects in human and animals [23].

Aflatoxins are produced by a polyketide pathway that pass through about twenty-seven enzymatic reactions which have been regulated by sets of genes including *nor-1*, *ver-1* and *omtA* have been shown to be involved in this process. *aflD* (*nor-1*) encodes a norsolorinic acid ketoreductase needed for the conversion of the 10-keto group of Norsolorinic Acid (NOR) to the 10-hydroxyl group of Versicolorin A (VERA) [24]. *aflM* (*ver-1*), predicted to encode a ketoreductase, is involved in the conversion of VERA into Sterigmatocystin (ST) [25]; *aflP* (*omtA*) codes for O-methyltransferase, which is one of the main genes responsible for transforming ST into O-Methylsterigmatocystin (OMST) that is the precursor for aflatoxin production [26].

Many other previous studies recorded detection of these genes in their *Aspergillus* isolates of food origin by various PCR techniques; Manonmani et al., [27], Rodrigues et al., [28] and Hassan et al., [29], who conducted several studies investigating the aflatoxigenicity of *Aspergillus* sp., could detect different genes in their *Aspergillus* isolates.

## Conclusion

It could be concluded that, breast samples revealed the highest contamination levels with *Aspergillus* sp.; in addition, *A. niger* was the prominently detected strain. PCR technique is a unique diagnostic tool for detection and identification of aflatoxigenic *Aspergillus* strains especially if the field of food safety. So, application of strict hygienic measures, proper use of water supply and food additives from good sources is recommended.

## References

1. Wong JT, Bruyn JD, Bagnol B, Grieve H, Li M, et al. Small-scale poultry and food security in resource-poor settings: A review. *Glob. Food Secur.* 2017; 15: 43-52.
2. Khalalfalla FA, Ali FHM and Saif-Alnasr MM. Microbiological quality of retail meats. *J. Vet. Med. Res.* 2017; 24: 311-321.
3. Leggieri MC, Toscano P and Battilani P. Predicted aflatoxin B1 increase in Europe due to climate change: actions and reactions at global level. *Toxins.* 2021; 13: 292.
4. Probst C, Njapau H and Cotty PJ. Outbreak of an acute aflatoxicosis in Kenya in 2004: identification of the causal agent. *Appl. Environ. Microbiol.* 2007; 73: 2762-2764.
5. Benkerroum N. Chronic and acute toxicities of aflatoxins: mechanisms of action. *Inter. J. Environ. Res. Pub. Hlth.* 2020; 17: 423.
6. Frisvad JC, Hubka V, Ezekiel CN, Hong SB, Nováková A, et al. Taxonomy of *Aspergillus* section Flavi and their production of aflatoxins, ochratoxins and other mycotoxins. *Studies in Mycology.* 2019; 93: 1-63.
7. Sohrabi N and Taghizadeh M. Molecular identification of aflatoxigenic *Aspergillus* species in feedstuff samples. *Curr. Med. Mycol.* 2018; 4: 1-6.
8. Sadhasivam S, Britzi M, Zakin V, Kostyukovsky M, Trostanetsky A, et al. Rapid detection and identification of mycotoxigenic fungi and mycotoxins in stored wheat grain. *Toxins.* 2017; 9: 302-310.
9. ISO. International Organization for Standardization. No.6887-1. Microbiology of the food chain- Preparation of test samples, initial suspension and decimal dilutions for microbiological examination-Part 1: General rules for the preparation of the initial suspension and decimal dilutions. 2017.
10. ISO. International Organization for Standardization. No.21527-1. Microbiology of food and animal feeding stuffs-Horizontal method for the enumeration of yeasts and moulds-Part 1: Colo-

- ny count technique in products with water activity greater than 0.95. 2008.
11. Pitt JI, Hocking, AD. *Fungi and Food Spoilage*. 2<sup>nd</sup> Ed, Springer, Boston, MA. 2009; 4: 19.
  12. Norlia M, Jinap S, Nor-Khaizura MAR, Radu S, Samsudin NIP et al. *Aspergillus* section *Flavi* and aflatoxins: occurrence, detection, and identification in raw peanuts and peanut-based products along the supply chain. *Front. Microbiol.* 2019; 10: 2602.
  13. Shaltout FA, Zakaria IM, Eltanani J and Elmelegy AS. Microbiological status of meat and chicken received to University student hostel. *Benha Vet. Med. J.* 2015; 29: 187-192.
  14. Lorenzo JM, Munekata PE, Dominguez R, Pateiro M, Saraiva JA, et al. Main groups of microorganisms of relevance for food safety and stability: general aspects and overall description. *Innov. Technol. Food Preserv.* 2018; 53-107.
  15. Benedict K, Chiller TM and Mody RK. Invasive fungal infections acquired from contaminated food or nutritional supplements: A review of the literature. *Foodborne Pathogens and Disease.* 2016; 13: 343-349.
  16. Richardson M and Rautemaa-Richardson R. Exposure to *Aspergillus* in home and healthcare facilities' water environments: focus on biofilms. *Microorganisms.* 2019; 7: 7-12.
  17. Darwish WS, El Bayomi RM, Abd El-Moaty AM and Gad TM. Mould contamination and aflatoxin residues in frozen chicken meat-cuts and giblets. *Jap. J. Vet. Res.* 2016; 64: 167-171.
  18. Shaltout FA, Nasief MZ, Lotfy LM and Gamil BT. Microbiological status of chicken cuts and its products. *Benha Vet. Med. J.* 2019; 37: 57-63.
  19. Hassan W. Some studies on effect of turmeric and lemon extracts on fungal contamination of chicken meat. Thesis, Master of Vet. Med. (Meat Hygiene), Benha Univ, Egypt. 2019.
  20. Abuzaid KE, Shaltout FA, Salem R and El-Diasty EM. Microbial aspect of some processed meat products with special reference to aflatoxins. *Benha Vet. Med. J.* 2020; 39: 24-28.
  21. OSHA" Occupational safety and Health Administration": A brief guide to mold in work place. OSHA Gov, California, USA. 2010.
  22. El-Abbasy TM. Mycological aspects of quail carcasses with a trial to improve their sanitary status. Thesis, Master of Vet. Med. (Meat Hygiene), Zagazig Univ, Egypt. 2007.
  23. Agriopoulou S, Stamatelopoulou E and Varzakas T. Advances in occurrence, importance, and mycotoxin control strategies: prevention and detoxification in Foods. *Foods (Basel, Switzerland).* 2020; 9: 137.
  24. Zhou R and Linz JE. Enzymatic function of the Nor-1 protein in aflatoxin biosynthesis in *Aspergillus parasiticus*. *Appl. Environ. Microbiol.* 1999; 65: 5639-5641.
  25. Henry KM and Townsend CA. Ordering the reductive and cytochrome P450 oxidative steps in demethylsterigmatocystin formation yields general insights into the biosynthesis of aflatoxin and related fungal metabolites. *J. Am. Chem. Soc.* 2005; 127: 3724-3733.
  26. Yabe K, Ando Y, Hashimoto J, Hamasaki T. Two distinct O-methyltransferases in Aflatoxin biosynthesis. *Appl. Environ. Microbiol.* 1989; 55: 2172-2177.
  27. Manonmani HK, Anand S, Chandrashekar A and Rati ER. Detection of aflatoxigenic fungi in selected food commodities by PCR. *Process Biochemistry.* 2005; 40: 2859-2864.
  28. Rodrigues P, Venencio A, Kozakiewicz Z and Lima N. A polyphasic approach to the identification of aflatoxigenic and non-aflatoxigenic strains of *Aspergillus* section *Flavi* isolated from Portuguese almonds. *Inter. J. Food Microbiol.* 2009; 129: 187-193.
  29. Hassan MN, El-Sayed AA and Nada HM. Detection of aflatoxins by HPLC and the expression of biosynthetic nor-1 gene of aflatoxin and ocrA gene of ochratoxin. *Benha Vet. Med. J.* 2015; 29: 1-10.