



# Large Right-Sided Complete Indirect Inguinal Hernia with Right Primary Vaginal Hydrocele

Shreya Soni<sup>1\*</sup>; Sheetal Asutkar<sup>1</sup>; Sandeep kumar Upadhyay<sup>2</sup>

<sup>1</sup>Department of Shalya Tantra, Mahatma Gandhi Ayurved College Hospital and Research Centre, Salod (H), Datta Meghe Institute of Higher Education and Research, Wardha 442001.

<sup>2</sup>Pg Scholar, Department of Shalya tantra, Datta meghe institute of Higher education and Research (DMIHER), Salod, Hirapur Wardha, India.

**\*Corresponding Author(s): Shreya Soni**

Department of Shalya Tantra, Mahatma Gandhi Ayurved College Hospital and Research Centre, Salod (H), Datta Meghe Institute of Higher Education and Research, Wardha 442001.

Email: dr.shreyasoni@gmail.com

Received: Jul 17, 2023

Accepted: Aug 01, 2023

Published Online: Aug 08, 2023

Journal: International Journal of Innovative Surgery

Publisher: MedDocs Publishers LLC

Online edition: <http://meddocsonline.org/>

Copyright: © Soni S (2023). This Article is distributed under the terms of Creative Commons Attribution 4.0 International License

**Keywords:** Zeiman's test; Irreducible; Inguinoscrotal; Pantaloon's; Abscess.

## Clinical Image

A male patient of 82 years old came, a Farmer by occupation came to the Outpatient of Shalya Tantra Department of Mahatma Gandhi Ayurved College and research centre with complaints of huge swelling in the Right Inguinoscrotal region which is irreducible in nature for 15 years. Initially, it was painless in nature as years passed dragging pain started and the swelling descended to the scrotum completely. Also, he had constipation for many years with increased frequency and intermittent urination. On Inspection-Size-Approx 30\*10cm (Length & Breadth), Cylindrical in shape, extending from above the inner part of the inguinal ligament down to the scrotum. The skin over the swelling is normal, the Impulse on coughing was positive, Penis is pushed towards another side. On Palpation-It was found to be Irreducible, Get above the swelling was negative,

Zeiman's test is positive on the Index finger, Elastic in consistency, Mild tenderness is elicited and the Left scrotum is found to be normal. On Percussion- a dull sound is elicited, Transillumination test positive. Chronic constipation, smoking, obesity, straining, and respiratory causes are the risk factors for hernia. Diagnostic criteria for Hernia with Hydrocele-Inguinal hernias are associated with hydrocele.

### Differential diagnosis

Groin abscess, Pantaloon's Hernia, Femoral hernia, Inguinal lymphadenitis, Scrotal oedema, Spermatocele.

### Final diagnosis

was found to be Right Indirect Complete Inguinal Hernia with Right Primary Hydrocele.



**Cite this article:** Soni S, Asutkar S, Sandeep kumar U. Large Right-Sided Complete Indirect Inguinal Hernia with Right Primary Vaginal Hydrocele. Int J Innov Surg. 2023; 6(2): 1035.

