



Anal buschke–lowenstein tumor

Mounia Bennani*; Jihane Ziani; Asmae Rasso; Sarra Elloudi; Zakia Douhi; Hanane Baybay; Fatima Zahra Mernissi
Hassan II hospital university, 81 lot riad avenue nasr fes, Morocco.

***Corresponding Author(s): Mounia Bennani**

Hassan II hospital university, 81 lot riad avenue nasr fes, Morocco.

Tel: 00212-6601-20505;

Email: mouniaabennani24@gmail.com

Received: Jul 22, 2019

Accepted: Sep 09, 2019

Published Online: Sep 12, 2019

Journal: Journal of Clinical Images

Publisher: MedDocs Publishers LLC

Online edition: <http://meddocsonline.org/>

Copyright: © Bennani M (2019). *This Article is distributed under the terms of Creative Commons Attribution 4.0 International License*

Clinical Image

Description

A 58-year-old patient, followed for a colorectal adenocarcinoma, initially placed under neoadjuvant radiotherapy, who developed 6 months later, an asymptomatic anal tumor gradually increasing in size.

The examination found: A budding tumor of 15 cm long axis extending from the anal margin, and taking all the inter-gluteal space, giving a hull crest appearance.

A cutaneous biopsy confirmed the diagnosis of lowenstein buschke tumor and the patient was referred to the visceral surgery department for possible tumor resection.

